

Land at Chepstow
Road
Raglan
Monmouthshire

**An Extended Phase 1
Habitat and Species
Assessment By:**






On Behalf Of:



September 2017

Client	Monmouthshire County Council
Project Name	Land at Chepstow Road, Raglan, Monmouthshire
Report Title	An Extended Phase 1 Habitat and Species Assessment
File Reference	GEN5513 (Revision 1)

Status	Signed	Name	Position	Date
Author		Carola Dallmeier BA (Hons) MSc Grad CIEEM	Ecologist	5 th August 2013
Author 1 st Revision		Carola Hoskins BSc (Hons) MSc ACIEEM	Senior Ecologist	5 th September 2017
Reviewer/Checker		Phillip L Morgan CEnv MCIEEM	Principal Ecologist	8 th September 2017

Members: Phillip Morgan CEnv MCIEEM Diane Morgan BA (Hons) ACIEEM
Carola Hoskins BA (Hons) MSc ACIEEM Robert Morgan
Issuing Office: Suite 131-136, Plas y Ffynnon, Cambrian Way, Brecon, LD3 7HP
Web Site: www.justmammals.co.uk **E-Mail:** carola@justmammals.co.uk **Telephone:** 01874 623616
Registered In: Cardiff **No:** OC345272 **Registered Address:** Llys Newydd, Llanfihangel Talyllyn, Brecon LD3 7TG
Vat No: 821 6376 35

Contents

	Page No.	
1	Executive Summary	1
2	Introduction	1
3	Surveyor Experience	2
4	Survey Methodology	2
5	Site Description	2
6	Survey Constraints	2
7	Survey Results	2 – 4
8	Discussion and Conclusions	4 – 5
9	Recommendations	5 – 7
10	References	8
	Appendix I: Site Location Plan	9
	Appendix II: Site Photographs	10
	Appendix III: Phase 1 Habitat Map	11
	Appendix IV: List of Recorded Species	12 – 13

List of Tables

Table	Description of Contents	Page No.
Table 1	Summary of Survey Activity and Weather Conditions	3
Table 2	Summary of Phase 1 Habitat Notes	3
Table 3	Recommended Native Tree and Shrub Species	6 – 7
Table 4	Recommended Garden Shrubs	7
Table 5	List of Recorded Species	12 – 13

List of Figures

Figure	Description of Contents	Page No.
Figure 1	Site location plan	9
Figure 2	Phase 1 habitat map	11

List of Plates

Plate	Description of Contents	Page No.
Plate 1	Looking west from entrance	10
Plate 2	Hedgerow along eastern boundary	10
Plate 3	View towards riparian corridor	10
Plate 4	Grassland and trees in south of property	10
Plate 5	Himalayan balsam along boundary	10
Plate 6	Water course	10

1 Executive Summary

- 1.1 An outline planning application for development of residential housing on land at Chepstow Road, Raglan, is being prepared by the present owners – Monmouthshire County Council. To support the proposals, an extended Phase 1 habitat survey was undertaken in July 2013, by an independent experienced ecologist. As no progress with the proposed development had occurred in the intervening period, an update survey was commissioned and carried out in August 2017.
- 1.2 The site is divided into two fields, both un-grazed, but mown at the time of the initial survey. A hedge is located along the eastern boundary, and the site abuts a small water course in the south. A total of 67 species of common and wide-spread plants were found, and the site is considered to be of medium ecological value. The update survey concluded that the site still holds the same ecological value in 2017. Careful design of any development will help retain some of the ecological value of the site, and appropriate recommendations are made in this report.
- 1.3 The riparian corridor at the southern end of the site is likely to be significant for commuting and feeding purposes for a number of species. Careful design will allow for its protection during and following the development.
- 1.4 Protected species and their potential presence on site were considered as part of the survey effort. Further assessment was originally recommended for otters and white clawed crayfish, and was subsequently undertaken (and covered in a separate report). A hedgerow along the western boundary is likely to be suitable for dormice, but additional survey is not recommended at this time as the low density presence of these animals is unlikely to be determined by traditional survey techniques in this instance. If the site is left undisturbed and undeveloped for much longer, it may also be necessary to carry out surveys for reptiles, due to the habitat steadily increasing in suitability for this group of animals. Small piles of debris are to be removed from site by hand, in case these are being used by animals for shelter.
- 1.5 Birds are likely to be nesting on site, therefore it is recommended to undertake any development work outside the breeding season for birds, between the months of March to August inclusive. Where this is not possible, an Ecological Clerk of Works will have to be in attendance when work is being carried out.
- 1.6 Himalayan balsam is widespread on the site, and the provision of a buffer zone throughout any development is recommended. Total eradication is unlikely to be achieved due to the high level of incursion by this plant on the surrounding land, but where undertaken, might free up the stream bank for amenity purposes.

2 Introduction

- 2.1 Planning permission is being sought for land off Chepstow Road, Raglan. Development will be for residential housing, the exact number of units not being defined at this point in time. As part of the site lies within the floodplain of the adjacent water course, some of the site will be left undeveloped, apart from a potential amenity or play area. The site, which is 2.08ha in size, lies at National Grid Reference (NGR) SO 4134 0738, at an altitude of approximately 37m Above Ordnance Datum.
- 2.2 The site is divided into two fields, and is unused. In order to assess the ecological value of the site and the potential for protected species to be present, the Just Mammals Consultancy LLP was commissioned to carry out an extended Phase 1 habitat assessment. A preliminary ecological appraisal/extended Phase 1 habitat survey was carried out in 2013, and an update survey was carried out in 2017, in order to assess whether the ecological conditions on site remained the same.
- 2.3 Different types of habitat were assessed, and the potential presence of protected species, such as badgers (*Meles meles*), bats, dormice (*Muscardinus avellanarius*), reptiles and amphibians, as well as nesting birds was considered. This report details the findings of the site assessment. Additionally, it makes recommendations concerning the ecological value of the site as well as the need for further survey work as appropriate.

3 Surveyor Experience

- 3.1 Surveyor, both for the original appraisal, and the latest update, was Carola Hoskins, who is also the author of this report. Carola holds an MSc in Environmental Conservation Management and has practical expertise with bats, birds, botanical assessments, mammalian and reptile surveys, both in Britain and overseas. As well as assisting in conservation-based research, she has carried out biodiversity audits and ecological enquiries. Carola has completed a study of water voles and is currently assisting with bird ringing. When Carola undertook the original survey she was a Graduate Member of the Chartered Institute of Ecology and Environmental Management (Grad CIEEM), but at the time of the August 2017, this had been upgraded to Associate Member (ACIEEM). Carola is a Senior Ecologist with the Just Mammals Consultancy LLP.

4 Survey Methodology

- 4.1 A botanical and habitat survey, and assessment for the presence and potential presence of protected species, was carried out on Tuesday the 30th of July 2013. The update survey was carried out on Wednesday the 16th of August 2017. Details of the survey activities and weather conditions are provided below in Table 1.
- 4.2 On both occasions, the site was walked over recording all plant species and features on a custom-made recording sheet. Habitats and notes were drawn onto a map of the survey site and digital camera photographs were taken. A coloured Phase 1 habitat map was produced, which can be found in Appendix III, and a list of plant species was recorded, as well as casual records of wildlife, which are shown in Appendix IV.
- 4.3 Assessment for the presence or potential presence of protected species, including bats, badgers, otters (*Lutra lutra*), dormice, reptiles and amphibians, was undertaken by considering the features of the site. Such features include grassland, hedgerows, buildings and trees. The potential suitability of the site for nesting birds was also considered.

5 Site Description

- 5.1 Situated on the outskirts of Raglan, along Chepstow Road, the site has a semi-urban character. It is made up of two fields of grassland; fencing suggests historic use for grazing animals. The site is level and covers a total area of just over two hectares. There is no evidence of previous development or usage for anything other than agricultural purposes, beyond the remnant base of a former large shed to the rear of the former Brooks Farm.
- 5.2 Boundaries of the site are mainly fences and hedgerows. The site is fenced off against housing (the former Brooks Farm), at the south-eastern edge. A fence/hedgerow runs along the boundary of the site with Chepstow Road, to the east. Where the site abuts the housing along Fayre Oaks, to the north, a fence runs along it, which continues along the western side of the property, where again housing is the adjacent land use. In the south, a small water course, the Nant y Wilcae, comprises the boundary of the site. A wire fence separates the two fields.
- 5.3 No part of the site is within a statutory designated site of conservation value (e.g. a Site of Special Scientific Interest (SSSI); Special Area of Conservation (SAC); Special Protection Area (SPA); or National Nature Reserve (NNR)). A search within a buffer zone of 2km around the site revealed no such designated sites to be in the area.

6 Survey Constraints

- 6.1 There were no constraints to the survey. Vegetation growth is mature. Access to the site was possible at all times both in 2013 and 2017.

7 Survey Results

- 7.1 Initial appraisal was undertaken on Tuesday the 30th of July 2013 by an experienced ecologist. Update survey was undertaken on Wednesday the 16th of August 2017, by the same ecologist. Details of the conditions under which survey was carried out are given in Table 1 below. Wind speeds given employ the Beaufort scale. Results of the survey efforts are separated by year.

Table 1: Summary of Survey Activity and Weather Conditions

Survey Type and Location	Dates	Timing	Weather Conditions
Day time visual inspection, botanical survey and habitat assessment, including protected species assessment	30/07/2013	10.50– 13:10 hours British Summer Time (BST)	Air temperature: 31.8°C Cloud cover: 4/8 oktas Wind speed: F2, light breeze Conditions: Dry
Day time visual inspection, botanical survey and habitat assessment, including protected species assessment	16/08/2017	12.00– 13:00 hours BST	Air temperature: 15°C Cloud cover: 5/8 oktas Wind speed: F2, light breeze Conditions: Dry
Surveyor	Carola Hoskins		

- 7.2 The site was divided into three different types of habitat for recording purposes. Table 2 below provides details of the various habitats and the species present in each of them.

Table 2: Summary of Phase 1 Habitat Notes

Habitat	Phase 1 Classification	Description of Area and Typical Species
Type 1	B2.2 Semi-improved neutral grassland	Both fields have a similar species composition with grasses making up most of the ground cover and some common forb species scattered in between. Dominant species are Cock's foot (<i>Dactylis glomerata</i>) and Yorkshire fog (<i>Holcus lanatus</i>). Also present are Perennial Ryegrass (<i>Lolium perenne</i>), creeping buttercup (<i>Ranunculus repens</i>) and pineappleweed (<i>Matricaria discoidea</i>). Target note: Some Himalayan balsam (<i>Impatiens glandulifera</i>) has spread from the water course into the grassland area.
Type 2	J2.1.1 Intact native, species-rich hedge	The hedge is made up of native species and intact. It runs along the eastern side of the property parallel with Chepstow Road and then along the boundary with the housing that was The Brooks Farm. It is approximately 1.5m wide and 2.5m high. Dominant species are hawthorn (<i>Crataegus monogyna</i>) and hazel (<i>Corylus avellana</i>). Also present are ash (<i>Fraxinus excelsior</i>) and bramble (<i>Rubus fruticosus</i>). Target note: Himalayan balsam has spread from the water course and is colonising along the hedgerow.
Type 3	A3.1 Scattered trees: tree line	Along the water course, the riparian corridor is mainly a tree line. The corridor is only about 5m wide; the trees are of a height of up to 10m. Ground vegetation is mainly of tall ruderal nature. Tree species present include alder (<i>Alnus glutinosa</i>) and willow (<i>Salix sp.</i>). Also present are meadowsweet (<i>Filipendula ulmaria</i>) and bramble. Target note: Himalayan balsam is present in large quantities along the riparian corridor and makes up the majority of the ground vegetation beneath the tree line. The invasion extends beyond the watercourse and beyond the site to the east and west.

- 7.3 A coloured Phase 1 habitat map was produced (see Appendix I). A total of 67 species of plants were recorded during the survey, as well as casual records of four birds and a number of other species. The site is considered to be of moderate ecological interest.
- 7.4 Both fields are recently mown to a short height of approximately 20cm. The arisings have been left on site. The hedgerow along the eastern end is of a height of approximately 2.5m, and interrupted in the middle by what was the former Brooks Farm, and is now housing. Connectivity is limited to the north where the main urban concentration of Raglan begins. To the south, the hedge connects to the riparian strip of woodland running along the southern end of the property, bordering the Nant y Wilcae.
- 7.5 Ecological assessment included consideration of the potential for protected species to be present on site. Bats use trees for roosting purposes and wooded areas for feeding purposes. None of the trees on site have any major cavities or rot holes conducive to usage by bats. There are no buildings on site that bats could be using for roosting purposes. It is likely that some foraging activity occurs over the site and that the riparian corridor is being used for commuting purposes.
- 7.6 Reptiles and amphibians use various kinds of habitat, including green spaces within urban surroundings. There are some potential foraging opportunities for reptiles on site; the watercourse along the southern end could potentially present hunting opportunities for grass snakes (*Natrix natrix*). There is little shelter available on site though and the close mowing of the

- grassland is likely to discourage reptiles from using the site. Two small piles of medium-sized rubble could potentially provide places of shelter for reptiles and amphibians.
- 7.7 Badgers will inhabit semi-urban agricultural land. Characteristic field signs of badgers are latrines dug into the ground where family groups deposit their dung, as well as badger setts, tunnel systems with multiple entrances. No such features were identified during the survey. Badgers, as a fairly large animal, also leave obvious runs in the vegetation. No runs were identified on site during the original survey, possibly owing to the recent mowing of the site. There is little north-bound connectivity and the road or water course would have to be crossed by any animals to gain access to the site, although this is entirely possible.
- 7.8 Hazel dormice make use of hedgerows where these are mature and provide sufficient foraging opportunity. Presence of these animals is indicated by nests found in the hedgerows and nuts opened in a way characteristic to the species. No evidence of dormice was found during the survey.
- 7.9 Otters inhabit riparian corridors of medium to large streams and will sometimes expand their territories onto smaller streams or use these for commuting purposes. No evidence of otters in form of footprints, spraints or holts, was found during this initial survey, although evidence for presence was established in a separate targeted survey (see separate report). The water course is relatively small and shallow along most of its length.
- 7.10 Water voles (*Arvicola amphibius*) can be found along slow-flowing small-medium-sized water courses, as well as lakes and ponds. They require dense ground vegetation and certain food species of plant, none of which are available at this site.
- 7.11 Consideration was made of the potential of the stream to be used by white clawed crayfish (*Austropotamobius pallipes*). As noted above the stream is relatively small and shallow, and in places there are deposits of silt. Whilst the banks are eroded and potential is limited for the species, it is possible that this species is present on the water course.
- 7.12 A large quantity of Himalayan balsam (*Impatiens glandulifera*) was found present on site in 2013. As is usual for this species, it colonises along the riparian corridor, pushing all other ground level vegetation out, and is now spreading along the hedgerow and into the field in places.
- 7.13 A number of birds were observed to be using the site during the survey. The hedgerow and tree line are both suitable for nesting purposes. The open fields were used for foraging purposes by insect-feeders. No nests were observed during the survey, but it is very likely the site is used for breeding purposes.
- 7.14 Update survey in 2017 revealed no significant changes to the site or its ecological value. Less Himalayan balsam was present on site than in 2013, but still occurs, especially along the water course. The spread into the grassland area is however considerably less, presumably an eradication effort has been carried out.
- 7.15 In terms of suitability for protected species, the site remains the same as it was in 2013. The suitability of the water course for otter and crayfish still remains the same, and an update survey was carried out at the same time, which is detailed in a separate report. The other ecological features of the site have not changed, the riparian habitat is an important corridor as per the above.

8 Discussion and Conclusions

- 8.1 The fields at Chepstow Road are of moderate interest to wildlife; the main grassland areas less so than the hedges. The riparian corridor offers some potential for wild animals and must not be disturbed as a consequence of any development, even acknowledging that both Chepstow Road, to the east, and the housing to the west, have already significantly encroached on the integrity of this corridor. As currently proposed, as a flood plain, the riparian corridor and its surroundings will be afforded some limited protection, within the development, which will also benefit wildlife.
- 8.2 Common and often ruderal species are present. The grassland is dominated by grass species and a limited diversity of herb species. Having recently been mown in 2013, the grassland held very limited interest for wildlife. Even so, a number of insects such as crickets and butterflies were observed, including comma (*Polygonia c-album*), and meadow brown (*Maniola jurtina*). During the 2017 survey the grass was significantly taller, although diversity of species remained

- largely the same. The hedges are mature and intact and provide foraging and commuting corridors for a number of species, including bats. The same holds true for the riparian corridor.
- 8.3 Protected species were considered during both surveys. There is no potential for the main part of the site to be used by bats for roosting. The riparian corridor and hedgerows are likely used for commuting purposes and feeding, some of the trees are suitable for roosting purposes. Any lighting scheme of the new development will have an impact on the usage of these corridors by bats as well as other species, such as otters.
- 8.4 No evidence, such as latrines or setts, to suggest the presence of badgers on the wider site was found during the surveys. However, the otter and crayfish survey both in 2014 and 2017 revealed a large badger sett to be present on the opposite bank of the stream, with well-established links to and presumably also through the stream. The site is therefore likely to be used by badgers for foraging purposes on a regular basis. The fields do not offer resting places for badgers, and as noted above, there are no setts in these areas.
- 8.5 The hazel dormouse is a species that has had considerable conservation effort directed towards it, yet the true distribution and size of the British is still somewhat unknown. The hedges and riparian trees bounding the site offer suitable habitat for these animals, and their presence cannot be dismissed. However, there is little woodland in the surrounding area – which would be an important reservoir for the species – so if they are present, it is likely to be at very low densities, and therefore difficult to determine.
- 8.6 Otters are now considered to be present along every water course in Wales. It is therefore highly likely that the Nant y Wilcae is inhabited by an otter or otters, which might be attracted along the water course to the several ponds which lie upstream – a likely source for amphibians in late winter/spring. In order to protect these animals the proposed buffer zone along the stream corridor must afford some protection and careful design of the development along with protective measures during construction will help in this process.
- 8.7 Water voles are making a recovery in parts of Wales where habitat is suitable. The Nant y Wilcae does not provide suitable habitat where it abuts the land at Chepstow Road. It is highly unlikely that water voles are present along this shaded water course.
- 8.8 There is moderate potential for the habitat to be used by reptiles. The small debris piles might have some hibernation potential. The riparian corridor might, in conjunction with the grassland habitat, provide suitable habitat for grass snakes. Since this will be largely undisturbed during the development, due to the floodplain zone to be specified, there is likely to be no significant impact on this species. However, if the habitat is left undisturbed for much longer, its suitability for reptiles will increase greatly, and colonisation by the more mobile species cannot be ruled out. Similarly, there is some small potential for native crayfish to be present. Again, provided the riparian zone is safeguarded during development, there is limited potential for these animals to be harmed.
- 8.9 Himalayan balsam is an invasive species and listed on Schedule 9 of the Wildlife and Countryside Act 1981 (as amended). As such, it is illegal to plant or otherwise cause to spread this invasive plant. Its distribution along the water course is such that the opposite bank to the property, as well as the areas up/downstream are all infested with this weed, making effective control or eradication highly unlikely.
- 8.10 A number of birds were observed during the survey and it is possible that some species are breeding on site. This will most likely be in the hedgerow or within the riparian tree line. Breeding efforts of all wild birds are protected under Schedule 1 of the Wildlife and Countryside Act 1981 (as amended).

9 Recommendations

- 9.1 Further survey is recommended for otters and white clawed crayfish. It is important to establish usage of the adjacent water course by otters in order to ascertain how these creatures use the rest of the site, and how best to proceed during the development in order to protect the otters from harm. Survey of both banks, looking for field signs such as spraints and holts, along the length of the site, as well as 250m either side of the proposed development, is recommended. Following survey, an ecological method statement can be formulated detailing how the otters will be protected during the development and where design of the new buildings and facilities will

- have to take otters into account, such as lighting schemes. Survey for otters can be carried out at any time of year.
- 9.2 A survey of the water course for white clawed crayfish is recommended. This protected species is increasingly rare on water courses, largely as a consequence of disease, but also as a consequence of development. White clawed crayfish can be surveyed for between July and early October.
- 9.3 Additional survey effort for dormice is unlikely to confirm their absence or otherwise. Dormouse nest tubes can be put up in the hedges at the site, but if dormice are present then it will be at very low densities and the chances of them finding or using the tubes is considered to be remote. It is therefore recommended that if a section of hedge must be removed as part of any development that it is inspected before hand to the presence of dormouse nests or any other field signs. The inspection must be carried out by an experienced ecologist and in the event that evidence for dormice presence is established work must be halted until a European Protected Species licence has been obtained from Natural Resources Wales. Where the development will not affect the hedgerows or disrupt connectivity with other habitat, no survey effort might be necessary. Optimum survey time for dormice is July to November.
- 9.4 Birds might be using the site for nesting purposes. It is therefore important that any work involving the hedgerow is carried out outside the nesting season from March to August inclusive. Where this is not possible, netting of the hedge outside the breeding season is possible in order to prevent nesting efforts being undertaken. Where neither of these options is valid, an Ecological Clerk of Works will have to be present during the works in order to search the hedgerow for nests. If an active nest be found, all work in the vicinity will have to cease until such a time when the breeding effort has been completed. Since no work is proposed along the riparian tree line, it is not expected that birds breeding in this habitat will be affected by the development.
- 9.5 Himalayan balsam on site will have to be a concern during and following the development. Due to its nature as an invasive plant, disturbance of it on a casual level must be prevented in order to avoid further spread of the plant. It is recommended to enforce a buffer zone around the infested area along the stream corridor during the development, going 5m beyond the furthest specimen. Depending on when the actual development starts, this might be quite far into the field, so localised control is advisable until such a time. The buffer zone must be separated from the development zone and no access for humans or machinery be allowed into the zone. The zone must not be used for storage or any other purposes. Control of the species in this area is unlikely to be efficient due to the surrounding areas being infested at the same level.
- 9.6 As a riparian corridor, the water course is vulnerable to pollution incidents caused by the development, another reason for the installation of a suitable buffer zone. The buffer zone will help keep any harmful substances away from the stream. Furthermore, night-time lighting of the site must be directed away from the stream corridor, so as to avoid disturbance of wildlife. This holds true for both during and after the development and careful lighting schemes will have to be devised.
- 9.7 It is considered unlikely that reptiles are using the site on a basis other than occasional and opportunistic at the present time. Consequently no further survey effort is recommended for this group. However, if the site continues to be left undisturbed for much longer, it is likely that it will be necessary to carry out survey for reptiles. As a precautionary measure, it is advisable to move the piles of debris found on site by hand in order to avoid damaging any creatures potentially using them for shelter. If reptiles or amphibians are found then advice must be sought from an ecologist immediately.
- 9.8 Development gives the opportunity to carry out enhancements to benefit wildlife. If the hedgerow(s) have to be removed as part of this proposal, replanting following development would be beneficial. Where possible, species used must be native. Table 3 below includes a list of suitable native species, which can be planted as part of the landscaping proposals. It is essential that such plants are sourced locally in order to reduce likelihood of importing diseases.

Table 3: Recommended Native Tree and Shrub Species

Common Name	Scientific Name
Alder	<i>Alnus glutinosa</i>
Crab apple	<i>Malus sylvestris</i>
Dogwood	<i>Cornus sanguinea</i>

Elder	<i>Sambucus nigra</i>
Field maple	<i>Acer campestre</i>
Hawthorn	<i>Crataegus monogyna</i>
Hazel	<i>Corylus avellana</i>
Holly	<i>Ilex aquifolium</i>
Rowan	<i>Sorbus aucuparia</i>
Silver birch	<i>Betula pendula</i>
Yew	<i>Taxus baccata</i>

- 9.9 Additional species which can be planted, which although not exclusively native species, will bring benefits for wildlife are included in Table 4 below. Again, only plants from local stockists must be used where this is possible.

Table 4: Recommended Garden Shrubs

Common Name	Scientific Name
Barberry	<i>Berberis vulgaris</i>
Clematis	<i>Clematis montana</i> or <i>Clematis vitalba</i>
Common broom	<i>Cytisus scoparius</i>
Dog rose	<i>Rosa canina</i>
Guelder rose	<i>Viburnum opulus</i>
Hebe	<i>Hebe albicans</i>
Honeysuckle	<i>Lonicera periclymenum</i>
Lavender	<i>Lavandula spp.</i>
Oregon grape	<i>Mahonia aquifolium</i>
Tree cotoneaster	<i>Cotoneaster 'Coral Beauty'</i>
Tree cotoneaster	<i>Cotoneaster Hybridus Pendulus</i>
Viburnum	<i>Viburnum davidii</i>


- 9.10 It is acceptable for other plant species to be provided on site, as recommended by the landscape architect. However, any planting proposals must include a minimum 70% proportion of the species listed in Tables 3 and 4.
- 9.11 Most developments include areas of grassland, and whilst some of these will require an amenity grassland seed mix, there are opportunities to sow wildflower grassland areas on parts of the site. To meet these needs it is recommended that the following seed mixes are used. British Seed Houses Mix A24 is a wear and tear mixture suitable for lawns and hard-working areas near to pathways. It contains five species of plant which are suitable for this location. For the wildflower areas the Emorsgate EM3 wildflower seed mix is recommended, with some twenty-five wild plant and grass species.
- 9.12 In order to benefit insects in particular, it is further recommended that additional seeding in the wildflower areas to encourage and benefit nectar feeding invertebrates, is carried out. An appropriate seed mix is available from Emorsgate – EN1. This mixture includes 23 plant species which can be added to the EM3 mix noted above.
- 9.13 It is important to implement good horticultural practice in any landscaping scheme, including the use of peat-free composts, mulches and soil conditioners. The use of pesticides (i.e. herbicides, insecticides, fungicides and slug pellets etc) must be discouraged to prevent cumulative fatal effects to animals via the food chain, particularly invertebrates, birds and/or mammals. Any pesticides used must be non-residual.
- 9.14 Ideally, as part of this proposal, the opportunity will be taken to develop a full and detailed biodiversity management plan. The plan would be expected to cover monitoring of all of the enhancement proposals set out above and the appointment of a biodiversity champion during and after the development.

10 References

- Gent, A.H., and Gibson, S.D. (Eds). (1998). '*Herpetofauna Worker's Manual*'. Joint Nature Conservation Committee, Peterborough
- Hayward, J. (1995). '*A New Key to Wild Flowers*'. University Press, Cambridge
- Hubbard, C.E. (1984). '*Grasses: A Guide to their Structure, Identification, Uses and Distribution in the British Isles*'. Penguin Books, London
- Inns, H. (2009). '*Britain's Reptiles and Amphibians*'. WildGuides Ltd., Old Basing
- Joint Nature Conservation Committee (1993). '*Handbook for Phase 1 Habitat Survey – A Technique for Environmental Audit*'. JNCC, Peterborough
- JNCC. (2004). '*Common Standards Monitoring Guidance for Reptiles and Amphibians*'. Joint Nature Conservation Committee, Peterborough
- Rose, F. (1981). '*The Wild Flower Key*'. Frederick Warne, London
- Streeter, D. (2009). '*Collins Flower Guide*'. Harper Collins, London
- Williams, C. (2010). '*Biodiversity for Low and Zero Carbon Buildings: A Technical Guide for New Build*'. RIBA Publishing, London

Appendix I: Site Location Plan

Figure 1: Site location plan



© Crown copyright and database rights (2017) Ordnance Survey (100055034)

Note: Survey boundary indicated in red

Appendix II: Site Photographs

Plate 1: Looking west from entrance



Plate 2: Hedgerow along eastern boundary



Plate 3: View towards riparian corridor



Plate 4: Grassland and trees in south of property



Plate 5: Himalayan balsam along boundary



Plate 6: Water course



Appendix III: Phase 1 Habitat Map

Figure 2: Phase 1 habitat map



Appendix IV: List of Recorded Species

Table 6: List of Recorded Species

Common Name	Scientific Name	1	2	3
Alder	<i>Alnus glutinosa</i>			•
Ash	<i>Fraxinus excelsior</i>		•	
Beech	<i>Fagus sylvatica</i>		•	
Bent grass, a	<i>Agrostis sp.</i>	•		
Black horehound	<i>Ballota nigra</i>	•		
Blackthorn	<i>Prunus spinosa</i>		•	
Bracken	<i>Pteridium aquilinum</i>	•		
Bramble	<i>Rubus fruticosus agg.</i>		•	
Broad-leaved dock	<i>Rumex obtusifolius</i>	•		
Charlock	<i>Sinapsis arvensis</i>		•	
Clover, a	<i>Trifolium sp.</i>	•		
Cock's-foot	<i>Dactylis glomerata</i>	•		
Colt's-foot	<i>Tussilago farfara</i>	•		
Common bird's-foot trefoil	<i>Lotus corniculatus</i>	•		
Common knotgrass	<i>Polygonum aviculare</i>	•		
Common nettle	<i>Urtica dioica</i>	•		
Common sorrel	<i>Rumex acetosa</i>	•		
Cow parsley	<i>Anthriscus sylvestris</i>	•		
Creeping buttercup	<i>Ranunculus repens</i>	•		
Dandelion	<i>Taraxacum agg.</i>	•		
Elder	<i>Sambucus nigra</i>		•	
False oat-grass	<i>Arrhenatherum elatius</i>	•		
Field bindweed	<i>Convolvulus arvensis</i>	•		
Fumitory	<i>Fumaria sp.</i>	•		
Good King Henry	<i>Chenopodium bonus henricus</i>	•		
Goosegrass	<i>Galium aparine</i>	•		•
Greater plantain	<i>Plantago major</i>	•		
Groundsel	<i>Senecio vulgaris</i>	•		
Hawthorn	<i>Crataegus monogyna</i>		•	•
Hazel	<i>Corylus avellana</i>		•	
Hedge bindweed	<i>Calystegia sepium</i>		•	
Hedge woundwort	<i>Stachys sylvatica</i>	•		
Herb-Robert	<i>Geranium robertianum</i>	•		
Himalayan balsam	<i>Impatiens glandulifera</i>	•	•	•
Hogweed	<i>Heracleum sphondylium</i>	•		
Holly	<i>Ilex aquifolium</i>		•	
Hop	<i>Humulus lupulus</i>		•	
Hop trefoil	<i>Trifolium campestre</i>	•		
Horsetail, a	<i>Equisetum spp.</i>	•		
Ivy	<i>Hedera helix</i>		•	
Lords and ladies	<i>Arum maculatum</i>	•		
Meadow buttercup	<i>Ranunculus acris</i>	•		
Meadow-grass, a	<i>Poa sp.</i>	•		
Meadowsweet	<i>Filipendula ulmaria</i>	•	•	•
Nipplewort	<i>Lapsana communis</i>	•		•
Oak	<i>Quercus sp.</i>	•		
Perennial rye-grass	<i>Lolium perenne</i>	•		
Pineappleweed	<i>Matricaria discoidea</i>	•		
Red clover	<i>Trifolium pratense</i>	•		
Redshank	<i>Persicaria maculosa</i>	•		
Ribwort plantain	<i>Plantago lanceolata</i>	•		
Rose, a	<i>Rose sp.</i>		•	
Rosebay willowherb	<i>Chamerion angustifolium</i>	•		
Rush, a	<i>Juncus sp.</i>	•		
Scarlet pimpernel	<i>Anagallis arvensis</i>	•		
Sedge, a	<i>Carex sp.</i>	•		
Sheep sorrel	<i>Rumex acetosella</i>	•		
Shepherd's-purse	<i>Capsella bursa-pastoris</i>	•		
Silverweed	<i>Potentilla anserina</i>	•		
Snow berry	<i>Symphoricarpos sp.</i>	•		
Speedwell, a	<i>Veronica sp.</i>	•		
Spurge, a	<i>Euphorbia sp.</i>	•		
Thistle	<i>Cirsium sp.</i>	•		•
Timothy	<i>Phleum pratense</i>	•		
White clover	<i>Trifolium repens</i>	•		
Willow, a	<i>Salix sp.</i>			•
Yarrow	<i>Achillea millefolium</i>	•		
Yorkshire fog	<i>Holcus lanatus</i>	•		

Fauna				
Green-veined white	<i>Pieris napi</i>			
Meadow brown (female)	<i>Maniola jurtina</i>			
Comma	<i>Polygonia c-album</i>			
Mole	<i>Talpa europaea</i>			
House martin	<i>Delichon urbicum</i>			
Raven	<i>Corvus corax</i>			
Magpie	<i>Pica pica</i>			
Blackbird	<i>Turdus merula</i>			

This Ecological Survey Report has been produced by the Just Mammals Consultancy LLP for Monmouthshire County Council solely. It may not be used by any person for any other purpose other than that specified without the express written permission of the Just Mammals Consultancy LLP. Any liability arising out of use by a third party of this document, for purposes not wholly connected with the above, shall be the responsibility of that party, who shall indemnify the Just Mammals Consultancy LLP against all claims, costs, damages, and losses, arising out of such use.

Unless specifically assigned or transferred within the terms of agreement, the consultant asserts and retains all Copyright, and other Intellectual Property Rights, in and over the Ecological Survey Report and its contents.

We confirm that in preparing this Ecological Survey Report we have exercised reasonable skill and care, taking into account the project objectives, the agreed scope of the work, prevailing site conditions, the degree of manpower and resources allocated to the project, and in compliance with the Chartered Institute of Ecology and Environmental Management's Code of Professional Conduct.

This Ecological Survey Report is valid for a period of two years from September 2017.

Just Mammals Consultancy is an ecological consultancy based in Mid-Wales. Specialising in legally protected mammal, reptile and amphibian species, but providing a wide range of ecological services, it provides appropriate expertise on behalf of a range of clients.

Clients include government departments, local and regional authorities, development agencies, commercial and industrial enterprises as well as statutory nature conservation organisations, wildlife trusts and other charitable bodies.

Please visit our web site www.justmammals.co.uk to see the full range of services we offer and some of the projects we have undertaken in the past.



Natural Problem Solvers

Just Mammals Consultancy LLP, Suite 131-136, Plas y Ffynnon, Cambrian Way, Brecon LD3 7HP Tel: 01874 623616